



Research Paper

## Central indices energy of special graphs

Saeed Mohammadian Semnani\*, Samira Sabeti

Department of Mathematics, Faculty of Mathematics, Statistics and Computer Science, Semnan University, Semnan, Iran

**Academic Editor:** Majid Arezoomand

**Abstract.** Given a graph  $G$  with vertex set  $V(G) = \{v_1, v_2, \dots, v_n\}$ . Let  $d_i$  be the degree of the vertex  $v_i$  in  $G$  for  $i = 1, 2, \dots, n$ . We introduce the sum of degrees and the product of degrees matrices of a graph. Furthermore, we consider the central indices matrix as an Arithmetic mean matrix, Geometric mean matrix, and Harmonic mean matrix. The spectral of these matrices has been computed. In this paper, we investigate the central indices energy of some classes of graphs and several results concerning its energy have been obtained.

**Keywords:** eigenvalue of a graph, energy, geometric mean energy, arithmetic mean energy, harmonic mean energy

**Mathematics Subject Classification (2010):** 05C50, 05C99.

### 1 Introduction

One of the most practical applications of graph theory to chemistry is in the creation of molecular descriptors. This kind of descriptor is based on invariants obtained from the representation of molecular structures like graphs. Most of these invariants are induced from the adjacency or distance matrices of the graphs. Significant classes of these descriptors are topological indices, which are simple numbers obtained by the mathematical methods of graphs associated with molecules. There are more than 120 topological indices but only at most a dozen have been applied. These can be classified by the structural properties of the

\*Email address: [S-mohammadian@semnan.ac.ir](mailto:S-mohammadian@semnan.ac.ir)

Received 27 May 2023; Revised 7 June 2023; Accepted 22 June 2023

First Publish Date: 1 July 2023

graphs used for their calculation [5].

In this paper we are concerned with simple finite graphs, without directed, multiple, or weighted edges, and without self-loops. Let  $A(G)$  be adjacency matrix of  $G$  and  $\lambda_1, \lambda_2, \dots, \lambda_n$  its eigenvalues. These are said to be the eigenvalues of the graph  $G$  and to form its spectrum. Graph energy is an invariant that is calculated from the eigenvalues of the adjacency matrix of the graph. In the mathematical literature, The concept of energy was formally put forward in 1978 [6]. Then, a dramatic change occurred, and graph energy started to attract the attention of a remarkably large number of mathematicians. The energy  $E(G)$  of the graph  $G$  is defined as the sum of the absolute values of its eigenvalues.

$$E(G) = \sum_{i=1}^n |\lambda_i|.$$

For more details on the mathematical aspects of the theory of graph energy see [1, 7, 16, 17]. Nowadays, due to the importance of applications of topological indices in chemistry, many researchers have studied these indices [8, 14]. Hence, one of the most well known is Wiener index [9, 18] is based on distance of vertices in the graph, the Hosoya index [12, 13] is calculated by counting non-incident edges in a graph, the energy and the Estrada index [10] are based on the spectrum of the graph, the Randic connectivity index [2, 11] and the Zagreb group indices [3, 15] are calculated using the degrees of vertices, geometric-arithmetic indices [4] is based on some properties of vertices of graph, etc.

Motivated by the mentioned papers, we study by focusing on the Central Indices energy of graphs. Plus, the energies of a number of well-known and much studied families of graphs are computed.

Let  $G = (V, E)$  be a simple graph of order  $n$  with vertex set  $V = V(G) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E = E(G)$ . Let  $d_i$  be the degree of the vertex  $v_i$  in  $G$  for  $i = 1, 2, \dots, n$ . The Sum of degrees matrix  $SD = SD(G)$  is

$$SD_{ij} = \begin{cases} d_i + d_j & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent} \\ 0 & \text{otherwise.} \end{cases}$$

The Arithmetic mean matrix  $AM = AM(G)$  is defined as

$$AM_{ij} = \begin{cases} \frac{d_i + d_j}{2} & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent} \\ 0 & \text{otherwise.} \end{cases}$$

The Product of degrees matrix  $PD = PD(G)$  is

$$PD_{ij} = \begin{cases} d_i d_j & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent} \\ 0 & \text{otherwise.} \end{cases}$$

The Geometric mean matrix  $GM = GM(G)$  is defined by

$$GM_{ij} = \begin{cases} \sqrt{d_i d_j} & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent} \\ 0 & \text{otherwise.} \end{cases}$$

Finally, the Harmonic mean matrix  $HM = HM(G)$  is defined by

$$HM_{ij} = \begin{cases} \frac{2d_i d_j}{d_i + d_j} & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent} \\ 0 & \text{otherwise.} \end{cases}$$

The above matrices are real symmetric, so we can order the eigenvalues of their matrices so that  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ . The Central Indices energy related to all above mention matrices of graph  $G$ , denoted by  $CIE(G)$ , is defined as

$$CIE(G) = \sum_{i=1}^n |\rho_i|.$$

## 2 Central indices energy of families of well-known graphs

In this section, we investigate the central indices energy of well-known of graphs. For  $n \geq 3$ , the Central indices energy of the Star graph  $S_n$  is as following table

Matrix name	Spectral	$E_{CI}(G)$
$A$	$\begin{pmatrix} \sqrt{n-1} & 0 & -\sqrt{n-1} \\ 1 & n-2 & 1 \end{pmatrix}$	$2\sqrt{n-1}$
$SD$	$\begin{pmatrix} n\sqrt{n-1} & 0 & -n\sqrt{n-1} \\ 1 & n-2 & 1 \end{pmatrix}$	$2n\sqrt{n-1}$
$AM$	$\begin{pmatrix} \frac{n\sqrt{n-1}}{2} & 0 & -\frac{n\sqrt{n-1}}{2} \\ 1 & n-2 & 1 \end{pmatrix}$	$n\sqrt{n-1}$
$PD$	$\begin{pmatrix} (n-1)\sqrt{n-1} & 0 & -(n-1)\sqrt{n-1} \\ 1 & n-2 & 1 \end{pmatrix}$	$2(n-1)\sqrt{n-1}$
$GM$	$\begin{pmatrix} n-1 & 0 & -(n-1) \\ 1 & n-2 & 1 \end{pmatrix}$	$2(n-1)$
$HM$	$\begin{pmatrix} \frac{2(n-1)\sqrt{n-1}}{n} & 0 & -\frac{2(n-1)\sqrt{n-1}}{n} \\ 1 & n-2 & 1 \end{pmatrix}$	$\kappa$

Where  $\kappa = \frac{4(n-1)\sqrt{n-1}}{n}$ .

*Proof.* We can see that the adjacency matrix of  $S_n$  as follow,

$$A(S_n) = \begin{pmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & 0 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \dots & 0 \end{pmatrix}$$

Also,  $SD = nA, AM = \frac{n}{2}A, PD = (n-1)A, GM = \sqrt{n-1}A, HM = \left(\frac{2(n-1)}{n}\right)A$ . With simple computation, we can get the results inside the table.  $\square$

We denote the complete graph on  $n$  vertices By  $K_n$ , and by  $\bar{K}_n$  its complement, i.e., the graph consisting of  $n$  isolated vertices. If  $G$  is the complete graph  $K_n$  then,

Matrix name	Spectral	$E_{CI}(G)$
A	$\begin{pmatrix} n-1 & -1 \\ 1 & n-1 \end{pmatrix}$	$2(n-1)$
SD	$\begin{pmatrix} 2(n-1)^2 & 2(n-1)^2 \\ 1 & n-1 \end{pmatrix}$	$4(n-1)^2$
AM	$\begin{pmatrix} (n-1)^2 - (n-1) \\ 1 & n-1 \end{pmatrix}$	$2(n-1)^2$
PD	$\begin{pmatrix} (n-1)^3 - (n-1)^2 \\ 1 & n-1 \end{pmatrix}$	$2(n-1)^3$
GM	$\begin{pmatrix} (n-1)^2 - (n-1) \\ 1 & n-1 \end{pmatrix}$	$2(n-1)^2$
HM	$\begin{pmatrix} (n-1)^2 - (n-1) \\ 1 & n-1 \end{pmatrix}$	$2(n-1)^2$

*Proof.* The Sum degree matrix of  $K_n$  is

$$SD(K_n) = \begin{pmatrix} 0 & 2(n-1) & 2(n-1) & \cdots & 2(n-1) \\ 2(n-1) & 0 & 2(n-1) & \cdots & 2(n-1) \\ 2(n-1) & 2(n-1) & 0 & \cdots & 2(n-1) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 2(n-1) & 2(n-1) & 2(n-1) & \cdots & 0 \end{pmatrix} = 2(n-1)(J_n - I_n)$$

where  $J_n$  is  $n \times n$  matrix with all entries 1 and  $I_n$  is the identity matrix. The characteristic polynomial of  $PD(K_n)$  is

$$\begin{aligned} f_m(G, \mu) &= \begin{vmatrix} \rho & -2(n-1) & -2(n-1) & \cdots & -2(n-1) \\ -2(n-1) & \rho & -2(n-1) & \cdots & -2(n-1) \\ -2(n-1) & -2(n-1) & \rho & \cdots & -2(n-1) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -2(n-1) & -2(n-1) & -2(n-1) & \cdots & \rho \end{vmatrix} \\ &= (\rho + 2(n-1)^2) \begin{vmatrix} \rho & -2(n-1) & -2(n-1) & \cdots & -2(n-1) \\ -1 & 1 & 0 & \cdots & 0 \\ -1 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -1 & 0 & 0 & \cdots & 1 \end{vmatrix} \\ &= (\rho + 2(n-1)^2)(\rho + 2(n-1))^{(n-1)} \end{aligned}$$

So,  $Spec_{SD}(K_n) = \begin{pmatrix} 2(n-1)^2 & -2(n-1) \\ 1 & n-1 \end{pmatrix}$ . So, Proof of calculation other central indices energy are similar to proof of Sum degree energy of complete graph. □

Let  $G$  be a graph on  $n$  vertices,  $n \geq 1$ . Then  $CIE(G) = 0$  if and only if  $G \cong \bar{K}_n$ . Let  $G = G_1 \cup G_2 \cup \dots \cup G_m$ . Then

(i)  $E(G) = E(G_1) + E(G_2) + \dots + E(G_m)$ .

(ii)  $GME(G) = GME(G_1) + GME(G_2) + \dots + GME(G_m)$ .

If the graph  $G$  is regular of degree  $k$ ,  $k > 0$ , then  $GME = kE(G)$ . If, in addition  $k = 0$ , then  $GME = 0$ .

*Proof.* If  $k = 0$ , then  $G \cong \bar{K}_n$ . From Lemma 2, we know that  $GME = 0$ . Suppose now that  $G$  is regular of degree  $k > 0$ , that is  $d_1 = d_2 = \dots = d_n = k$ . Then all non-zero terms in  $GM$  matrix are equal to  $k$ . This implies that  $GM(G) = kA(G)$ . Then we have  $\rho_i = k\lambda_i$ , and therefore  $GME = kE(G)$ . □

If  $G$  is the complete bipartite graph  $K_{n,n}$ , the spectral and energy of central indices are calculated as follow,

Matrix name	Spectral	$E_{CI}(G)$
A	$\begin{pmatrix} n & 0 & -n \\ 1 & 2n-2 & 1 \end{pmatrix}$	$2n$
SD	$\begin{pmatrix} 2n^2 & 0 & -2n^2 \\ 1 & 2n-2 & 1 \end{pmatrix}$	$4n^2$
AM	$\begin{pmatrix} n^2 & 0 & -n^2 \\ 1 & 2n-2 & 1 \end{pmatrix}$	$2n^2$
PD	$\begin{pmatrix} n^3 & 0 & -n^3 \\ 1 & 2n-2 & 1 \end{pmatrix}$	$2n^3$
GM	$\begin{pmatrix} n^2 & 0 & -n^2 \\ 1 & 2n-2 & 1 \end{pmatrix}$	$2n^2$
HM	$\begin{pmatrix} n^2 & 0 & -n^2 \\ 1 & 2n-2 & 1 \end{pmatrix}$	$2n^2$

*Proof.* The proof is similar to Theorem 2. □

If  $G$  is the complete bipartite graph  $K_{n,m}$ , with  $n + m$  vertices. the spectral and energy of

central indices are calculated as follow,

Matrix name	Spectral	$E_{CI}(G)$
A	$\begin{pmatrix} \sqrt{mn} & 0 & -\sqrt{mn} \\ 1 & m+n-2 & 1 \end{pmatrix}$	$2\sqrt{mn}$
SD	$\begin{pmatrix} (m+n)\sqrt{mn} & 0 & -(m+n)\sqrt{mn} \\ 1 & m+n-2 & 1 \end{pmatrix}$	$2(m+n)\sqrt{mn}$
AM	$\begin{pmatrix} \frac{(m+n)\sqrt{mn}}{2} & 0 & -\frac{(m+n)\sqrt{mn}}{2} \\ 1 & m+n-2 & 1 \end{pmatrix}$	$(m+n)\sqrt{mn}$
PD	$\begin{pmatrix} mn\sqrt{mn} & 0 & -mn\sqrt{mn} \\ 1 & m+n-2 & 1 \end{pmatrix}$	$2mn\sqrt{mn}$
GM	$\begin{pmatrix} mn & 0 & -mn \\ 1 & m+n-2 & 1 \end{pmatrix}$	$2mn$
HM	$\begin{pmatrix} \frac{2mn\sqrt{mn}}{m+n} & 0 & -\frac{2mn\sqrt{mn}}{m+n} \\ 1 & m+n-2 & 1 \end{pmatrix}$	$\kappa'$

Where  $\kappa' = \frac{4mn\sqrt{mn}}{m+n}$ .

*Proof.* The proof is similar to Theorem 2. □

Let  $G$  be a bipartite complete graph, we know number of edges equal  $mn$  then the Geometric mean energy is twice the number of edges, that is  $GME = 2mn$ . The crown graph  $S_n^0$  for an integer  $n \geq 3$  is the graph with vertex set  $\{u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n\}$  and edge set  $\{u_i v_j : 1 \leq i, j \leq n, i \neq j\}$ . Therefore  $S_n^0$  coincides with the complete bipartite graph  $K_{n,n}$  with the horizontal edges removed. If  $G$  is the crown graph  $S_n^0$  with  $2n$  vertices, the Central Indices energy is defined

Matrix name	Spectral	$E_{CI}(G)$
A	$\begin{pmatrix} n-1 & 1 & -1 & -(n-1) \\ 1 & n-1 & n-1 & 1 \end{pmatrix}$	$2n$
SD	$\begin{pmatrix} 2(n-1)^2 & 2(n-1) & -2(n-1) & -2(n-1)^2 \\ 1 & n-1 & n-1 & 1 \end{pmatrix}$	$8(n-1)^2$
AM	$\begin{pmatrix} (n-1)^2 & n-1 & -(n-1) & -(n-1)^2 \\ 1 & n-1 & n-1 & 1 \end{pmatrix}$	$4(n-1)^2$
PD	$\begin{pmatrix} (n-1)^3 & (n-1)^2 & -(n-1)^2 & -(n-1)^3 \\ 1 & n-1 & n-1 & 1 \end{pmatrix}$	$4(n-1)^3$
GM	$\begin{pmatrix} (n-1)^2 & n-1 & -(n-1) & -(n-1)^2 \\ 1 & n-1 & n-1 & 1 \end{pmatrix}$	$4(n-1)^2$
HM	$\begin{pmatrix} (n-1)^2 & n-1 & -(n-1) & -(n-1)^2 \\ 1 & n-1 & n-1 & 1 \end{pmatrix}$	$4(n-1)^2$

If the graph  $G$  is regular of degree  $k, k > 0$ , then  $PDE = k^2E(G)$ . If, in addition  $k = 0$ , then  $PDE = 0$ .

*Proof.* If  $k = 0$ , then  $G \cong \bar{K}_n$ . From Lemma 2, we know that  $PDE = 0$ . Suppose now that  $G$  is regular of degree  $k > 0$ , that is  $d_1 = d_2 = \dots = d_n = k$ . Then all non-zero terms in  $GM$  matrix are equal to  $k$ . This implies that  $PD(G) = k^2 A(G)$ . Then we have  $\rho_i = k^2 \lambda_i$ , and therefore  $PDE = kE(G)$ .  $\square$

If the graph  $G$  is regular of degree  $k$ ,  $k > 0$ , then

$$AME(G) = GME(G) = HME(G).$$


## Acknowledgements

This research was supported by Research Council of Semnan University.

## References

- [1] R. B.apat, Graphs and matrices, New York, Springer, 27, 2010.
- [2] B. Borovicanin, E. Zogic, On Randic energy of a graph, In The 14th Serbian Mathematical Congress (14th SMAK), Kragujevac, 2018.
- [3] B. Borovicanin, T.A. Lampert, On the maximum and minimum Zagreb indices of trees with a given number of vertices of maximum degree, MATCH Commun. Math. Comput. Chem. 74 (2015) 81-96.
- [4] K. C. Das, I. Gutman, B. Furtula, Survey on geometric-arithmetic indices of graphs, MATCH Commun. Math. Comput. Chem. 65(3) (2011) 595-644.
- [5] E. Estrada, A. Ramírez, Edge adjacency relationships and molecular topographic descriptors. Definition and QSAR applications, J. Chem. Inf. Comput. Sci. 36(4) (1996) 837-843.
- [6] I. Gutman, The energy of a graph: old and new results, In: Algebraic combinatorics and applications, Springer, Berlin, Heidelberg, 1987.
- [7] I. Gutman, B. Furtula, E. Zogic, E. Glogic, Resolvent energy of graphs, MATCH Commun. Math. Comput. Chem. 75 (2016) 279-290.
- [8] I. Gutman, B. Furtula (Eds.), Novel Molecular Structure Descriptors-Theory and Applications I , Univ. Kragujevac, Kragujevac, 2010.
- [9] I. Gutman, S. Klavzar, B. Mohar (Eds.), Fifty years of the Wiener index, MATCH Commun. Math. Comput. Chem. 35 (1997) 1-259.
- [10] A. Jahanbani, H.H. Raz, On the harmonic energy and Estrada index of graphs, MATI. 1(1) (2019) 1-20.
- [11] H. Jun, Y.M. Liu, J.K. Tian, Note on the Randic energy of graphs, Kragujev. J. Math. 42(2) (2018) 209-215.
- [12] S. Li, X. Li, Z. Zhu, On minimal energy and Hosoya index of unicyclic graphs, MATCH Commun. Math. Comput. Chem. 61 (2009) 325-339.
- [13] P. D. Manuel, An efficient Hosoya index algorithm and its application, IJCAET. 11(2) (2019) 196-205.
- [14] O. Mekenyan, D. Bonchev, N. Trinajstic, Chemical graph theory: Modeling the thermodynamic properties of molecules, Int. J. Quantum Chem. 18(2) (1980) 369-380.
- [15] N.J. Rad, A. Jahanbani, I. Gutman, Zagreb energy and Zagreb estrada index of graphs, Match. Commun. Math. Comput. Chem. 79 (2018) 371-386.
- [16] S. Sabeti, A. B. Dehkordi, S. M. Semnani, The minimum edge covering energy of a graph. Kragujev. J. Math. 45(6) (2021) 969-975.
- [17] S. K. Vaidya, K. M. Popat, Some new results on energy of graphs, MATCH Commun. Math. Comput. Chem. 77 (2017) 589-594.
- [18] X. D. Zhang, Y. Liu, M. X. Han, Maximum Wiener index of trees with given degree sequence, MATCH Commun. Math. Comput. Chem. 64 (2010) 661-682.

**Citation:** S. Mohammadian Semnani, S. Sabeti, Central indices energy of special graphs, J. Disc. Math. Appl. 8(2) (2023) 105–112.

 <https://doi.org/10.22061/jdma.2023.1640>



**COPYRIGHTS**

©2023 The author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.