



Research Paper

On the Covering Radius of DNA Code over a Finite Ring

P. Chella Pandian*

Department of Mathematics, Srimad Andavan Arts and Science College(A), Tiruchirappalli,
Taminadu, India

Academic Editor: Modjtaba Ghorbani

Abstract. In this paper, lower bound and upper bound on the covering radius of DNA codes over a finite ring \mathbb{N} with respect to chinese euclidean distance are given. Also determine the covering radius of various Repetition DNA codes, Simplex DNA code Type α and Simplex DNA code Type β and bounds on the covering radius for MacDonal DNA codes of both types over \mathbb{N} .

Keywords: DNA Code, Finite Ring, Covering Radius, Simplex Codes

Mathematics Subject Classification (2010): 94B25, 94B05, 11H31, 11H71.

1 Introduction

In, DNA is found naturally as a double stranded molecule, with a form similar to a twisted ladder. The backbone of the DNA helix is an alternating chain of sugars and phosphates, while the association between the two strands is variant combinations of the four nitrogenous bases adenine (A), thymine (T), guanine (G) and cytosine (C). The two ends of the strand are distinct and are conventionally denoted as 3' end and 5' end. Two strands of DNA can form (under suitable conditions) a double strand if the respective bases are Watson- Crick [17] complements of each other - A matches with T and C matches with G, also 3' end matches with 5' end.

The problem of designing DNA codes (sets of words of fixed length n over the alphabets $\{A, C, G, T\}$) that satisfy certain combinatorial constraints has applications for reliably storing and retrieving information in synthetic DNA strands. These codes can be used in

*Email address: chellapandianpc@gmail.com, chella@andavancollege.ac.in

Received 26 April 2023; Revised 4 May 2023; Accepted 22 May 2023

First Publish Date: 1 June 2023

particular for DNA computing [1] or as molecular bar-codes.

There are many researchers doing research on code over finite rings. In particular, codes over \mathbb{Z}_4 received much attention [2–5,9,11,15,16]. The covering radius of binary linear codes were studied [4,5]. Recently the covering radius of codes over \mathbb{Z}_4 has been investigated with respect to chinese euclidean distances [14]. In 1999, Sole et al gave many upper and lower bounds on the covering radius of a code over \mathbb{Z}_4 with chinese euclidean distances. In [5,13], the covering radius of some particular codes over \mathbb{Z}_4 have been investigated.

In this correspondence, consider a finite ring \mathbb{N} . In this paper, Investigate the covering radius of the Simplex DNA codes of both types and MacDonald DNA codes and repetition DNA codes over \mathbb{N} . Also generalized some of the known bounds in [2].

2 Preliminary

Coding theory has several applications in Genetics and Bioengineering. The problem of designing DNA codes (sets of that words of fixed length n over the alphabet $\mathbb{N} = \{A, C, G, T\}$ that satisfy certain combinatorial constraints) has applications for reliably storing and retrieving information in synthetic DNA strands.

A DNA code of length n is a set of codewords (x_1, x_2, \dots, x_n) with $x_i \in \{A, C, G, T\} = \mathbb{N}$ (representing the four nucleotides in DNA). Use a hat to denote the Watson-Crick complement of a nucleotide, so A matches with T and C matches with G .

The DNA codes are sets of words of fixed length n over the alphabet $\mathbb{N} = \{A, C, G, T\}$ and it follows the map $A \rightarrow 0, C \rightarrow 1, T \rightarrow 2$ and $G \rightarrow 3$. Therefore the problem of the DNA codes is corresponding to the problem of the \mathbb{Z}_4 -linear codes. These transpositions do not affect the GC-weight of the codeword (the number of entries that are C or G). In my work, by using the above map in \mathbb{Z}_4 with chinese euclidean weight, so obtain the covering radius for repetition DNA codes .

Let $d = (d_1, d_2, \dots, d_n) \in \mathbb{N}^n$ and n be its length. Let b be an element of $\{A, C, G, T\}$. For all $d = (d_1, d_2, \dots, d_n) \in \mathbb{N}^n$, define the weight of d at b to be $w_b(d) = |\{i | x_i = b\}|$.

A DNA linear code C of length n over \mathbb{N} is an additive subgroup of \mathbb{N}^n . An element of C is called a DNA codeword of C and a generator matrix of C is a matrix whose rows generate C .

In [14], the chinese Euclidean weight $w(x)$ of a vector x is $\sum_{i=1}^n \left\{ 2 - 2 \cos \left(\frac{2\pi x_i}{4} \right) \right\}$.

A linear Gray map φ from $\mathbb{N}_4 \rightarrow \mathbb{Z}_2^2$ is defined by $\varphi(x + 2y) = (y, x + y)$, for all $x + 2y \in \mathbb{N}$. The image $\varphi(C)$, of a linear code C over \mathbb{N} of length n by the Gray map is a binary code of length $2n$ with same cardinality [15].

Any DNA linear code C over \mathbb{N} is equivalent to a code with Generator Matrix(GM) of the form

$$GM = \begin{bmatrix} I_{k_0} & A & B \\ 0 & 2I_{k_1} & 2D \end{bmatrix}, \text{ where } A, B \text{ and } D \text{ are matrices over } \mathbb{N}.$$

Then the DNA code C contain all DNA codewords $[v_0, v_1]GM$, where v_0 is a vector of length k_1 over \mathbb{N} and v_1 is a vector of length k_2 over \mathbb{Z}_2 . Thus C contains a total of $4^{k_1}2^{k_2}$

codewords. The parameters of C are given $[n, 4^{k_1} 2^{k_2}, d]$, where d represents the minimum chinese Euclidean distance of C .

A DNA linear code C over \mathbb{N} of length n , 2-dimension k , minimum chinese euclidean distance d is called an $[n, k, d_{CE}]$ or simply an $[n, k, d]$ code.

In this paper, define the covering radius of dna codes over \mathbb{N} with respect to chinese euclidean distance and in particular study the covering radius of Simplex DNA codes of type α and type β namely, S_k^α and S_k^β and their MacDonald DNA codes and repetition DNA codes over \mathbb{N} . Section 2 contains basic results for the covering radius of DNA codes over \mathbb{N} . Section 3 determines the covering radius of different types of repetition DNA codes. Section 4 determines the covering radius of Simplex DNA codes and finally section 5 determines the bounds on the covering radius of MacDonald DNA codes.

3 Covering radius of repetition DNA codes

Let d be a chinese euclidean distance a DNA code C over \mathbb{N} . Thus, the covering radius of C :

$$r_d(C) = \max_{u \in \mathbb{N}^n} \left\{ \min_{c \in C} \{d(c, u)\} \right\}.$$

The following result of Mattson [6] is useful for computing covering radius of codes over rings generalized easily from codes over finite fields.

If C_0 and C_1 are codes over \mathbb{N} generated by matrices GM_0 and GM_1 respectively and if C is the code generated by $GM = \left(\begin{array}{c|c} 0 & GM_1 \\ \hline GM_0 & A \end{array} \right)$ then $r_d(C) \leq r_d(C_0) + r_d(C_1)$ and the covering radius of D (concatenation of C_0 and C_1) satisfy the following $r_d(D) \geq r_d(C_0) + r_d(C_1)$, for all distances d over \mathbb{N} .

A q -ary repetition code C over a finite field $\mathbb{F}_q = \{\alpha_0 = 0, \alpha_1 = 1, \alpha_2, \alpha_3, \dots, \alpha_{q-1}\}$ is an $[n, 1, n]$ code $C = \{\bar{\alpha} \mid \alpha \in \mathbb{F}_q\}$, where $\bar{\alpha} = (\alpha, \alpha, \dots, \alpha)$. The covering radius of C is $\left\lceil \frac{n(q-1)}{q} \right\rceil$ [11]. Using this, it can be seen easily that the covering radius of block of size n repetition code $[n(q-1), 1, n(q-1)]$ generated by

$$GM = \left[\overbrace{11 \dots 1}^n \overbrace{\alpha_2 \alpha_2 \dots \alpha_2}^n \overbrace{\alpha_3 \alpha_3 \dots \alpha_3}^n \dots \overbrace{\alpha_{q-1} \alpha_{q-1} \dots \alpha_{q-1}}^n \right]$$

is $\left\lceil \frac{n(q-1)^2}{q} \right\rceil$, since it will be equivalent to a repetition code of length $(q-1)n$.

Consider the repetition dna code over \mathbb{N} . There are two types of them of length n viz.

- cytosine repetition code $C_\beta : [n, 1, 2n]$ generated by $GM_\beta = \left[\overbrace{C \ C \dots C}^n \right]$
- thymine repetition code $C_\alpha : (n, 2, 4n)$ generated by $GM_\alpha = \left[\overbrace{T \ T \dots T}^n \right]$.

Theorem 3.1. Let C_β and C_α be the dna code of type β and α type in generator matrices GM_β and GM_α . Then, $4 \left\lfloor \frac{n}{2} \right\rfloor \leq r(C_\alpha) \leq 2n$ and $r(C_\beta) = 2n$.

Proof. Let $x = \overbrace{TT \cdots T}^{\lfloor \frac{n}{2} \rfloor} \overbrace{AA \cdots A}^{\lceil \frac{n}{2} \rceil}$ and the code of $C = \{AA \cdots A, TT \cdots T\}$ is generated by $[TT \cdots T]$ is an $[n, 1, 2n]$ code. Then, $d(x, AA \cdots A) = wt(x - AA \cdots A) = 4 \lceil \frac{n}{2} \rceil$ and $d(x, TT \cdots T) = wt(x - TT \cdots T) = 4 \lfloor \frac{n}{2} \rfloor$. Therefore $d(x, C_\alpha) = \min \{4 \lceil \frac{n}{2} \rceil, 4 \lfloor \frac{n}{2} \rfloor\}$. Thus, by definition of covering radius

$$r(C_\alpha) \geq 4 \lfloor \frac{n}{2} \rfloor \tag{1}$$

Let x be any word in \mathbb{N}^n . Let us take x has ω_0 coordinates as 0's, ω_1 coordinates as 1's, ω_2 coordinates as 2's, ω_3 coordinates as 3's, then $\omega_0 + \omega_1 + \omega_2 + \omega_3 = n$. Since $C_\alpha = \{AA \cdots A, TT \cdots T\}$ and chinese euclidean weight of $\mathbb{N} : 0$ is 0, C and G is 2 and T is 4. Therefore, $d(x, 00 \cdots 0) = n - \omega_0 + \omega_1 + 3\omega_2 + \omega_3$ and $d(x, TT \cdots T) = n - \omega_2 + \omega_1 + 3\omega_0 + \omega_3$.

Thus $d(x, C_\alpha) = \min \{n - \omega_0 + \omega_1 + 3\omega_2 + \omega_3, n - \omega_2 + \omega_1 + 3\omega_0 + \omega_3\}$ and hence,

$$d(x, C_\alpha) \leq n + n = 2n. \tag{2}$$

Hence, from the Equation (1) and (2), so $4 \lfloor \frac{n}{2} \rfloor \leq r(C_\alpha) \leq 2n$.

Now, obtain the covering radius of C_β covering with respect to the chinese euclidean weight. Then $d(x, AA \cdots A) = n - \omega_0 + \omega_1 + 3\omega_2 + \omega_3, d(x, CC \cdots C) = n - \omega_1 + \omega_0 + \omega_2 + 3\omega_3, d(x, TT \cdots T) = n - \omega_2 + 3\omega_1 + \omega_3$ and $d(x, GG \cdots G) = n - \omega_3 + 3\omega_1 + \omega_0 + \omega_2$, for any $x \in \mathbb{N}^n$.

This implies $d(x, C_\beta) = \min \{n - \omega_0 + \omega_1 + 3\omega_2 + \omega_3, n - \omega_1 + \omega_0 + \omega_2 + 3\omega_3, n - \omega_2 + 3\omega_1 + \omega_3, n - \omega_3 + 3\omega_1 + \omega_0 + \omega_2\} \leq 2n$ and hence $r(C_\beta) \leq 2n$.

Let $x = \overbrace{AA \cdots A}^t \overbrace{CC \cdots C}^t \overbrace{TT \cdots T}^t \overbrace{GG \cdots G}^{n-3t}$, where $t = \lfloor \frac{n}{4} \rfloor$, then $d(x, AA \cdots A) = 2n, d(x, CC \cdots C) = 4n - 8t, d(x, TT \cdots T) = 2n$ and $d(x, GG \cdots G) = 8t$.

Therefore $r(C_\beta) \geq \min \{2n, 4n - 8t, 8t\} \geq 2n$. □

Block repetition code

Let $GM = \overbrace{[CC \cdots C]}^n \overbrace{[TT \cdots T]}^n \overbrace{[GG \cdots G]}^n$ be a generator matrix of \mathbb{N} in each block of repetition code length is n . Then, the parametrs of Block Repetition Code(BRC) is $[3n, 1, 8n]$. The code of $BRC = \{c_0 = A \cdots AA \cdots AA \cdots A, c_1 = C \cdots CT \cdots TG \cdots G, c_2 = T \cdots TA \cdots AT \cdots T, c_3 = G \cdots GT \cdots TC \cdots C\}$, dimension of BRC is 1 and chinese euclideran weight is $8n$. Note that, the block repetition code has constant chinese euclidean weight is $8n$.

Theorem 3.2. To find $4 \lfloor \frac{n}{2} \rfloor + 4n \leq r(BRC^{3n}) \leq 6n$.

Proof. Let $x = AA \cdots A \in \mathbb{N}^{3n}$. Then, $d(x, BRC^{3n}) = 4 \lfloor \frac{n}{2} \rfloor + 4n$. Hence by definition, $r(BRC^{3n}) \geq 4 \lfloor \frac{n}{2} \rfloor + 4n$.

Let $x = (u|v|w) \in \mathbb{N}^{3n}$ with u, v and w have compositions (r_0, r_1, r_2, r_3) , (s_0, s_1, s_2, s_3) and (t_0, t_1, t_2, t_3) respectively such that $\sum_{i=0}^3 r_i = n, \sum_{i=0}^3 s_i = n$ and $\sum_{i=0}^3 t_i = n$, then

$d(x, c_0) = 3n - r_0 + r_1 + 3r_2 + r_3 - s_0 + s_1 + 3s_2 + s_3 - t_0 + t_1 + 3t_2 + t_3, d(x, c_1) = 3n - r_1 + r_0 + r_2 + 3r_3 - s_2 + 3s_0 + s_1 + s_0 - t_3 + t_0 + 3t_1 + t_2, d(x, c_2) = 3n - r_2 + r_1 + 3r_0 + r_3 - s_0 + s_1 + 3s_2 + s_3 - t_2 + 3t_0 + t_1 + t_3$ and $d(x, c_3) = 3n - r_3 + 3r_1 + r_0 + r_2 - s_2 + 3s_0 + s_1 + s_3 - t_1 + 3t_3 + t_0 + t_2$.

Thus, $d(x, BRC) = \min\{3n - r_0 + r_1 + 3r_2 + r_3 - s_0 + s_1 + 3s_2 + s_3 - t_0 + t_1 + 3t_2 + t_3, 3n - r_1 + r_0 + r_2 + 3r_3 - s_2 + 3s_0 + s_1 + s_3 - t_3 + t_0 + 3t_1 + t_2, 3n - r_2 + r_1 + 3r_0 + r_3 - s_0 + s_1 + 3s_2 + s_3 - t_2 + 3t_0 + t_1 + t_3, 3n - r_3 + 3r_1 + r_0 + r_2 - s_2 + 3s_0 + s_1 + s_3 - t_1 + 3t_3 + t_0 + t_2\} \leq 6n$ and hence, $r(BRC^{3n}) \leq 6n$. \square

Define a two block repetition dna code over \mathbb{N} of each of length is n and the parameters of two block repetition cod $BRC^{2n} : [2n, 1, 4n]$ is generated by $G = [\overbrace{CC \cdots C}^n \overbrace{TT \cdots T}^n]$. Use the above and obtain a following

Theorem 3.3. $4 \lfloor \frac{n}{2} \rfloor + 2n \leq r(BRC^{2n}) \leq 4n$.

Let $GM = [\overbrace{C C \cdots C}^m \overbrace{T T \cdots T}^n]$ be the generalized generator matrix for two different block repetition dna code of length are m and n respectively. In the parameters of two different block repetition code (BRC^{m+n}) are $[m + n, 1, \min\{4m, 3m + 3n\}]$ and Theorem 3.3 can be easily generalized for two different length using similar arguments to the following.

Theorem 3.4. $2m + 4 \lfloor \frac{n}{2} \rfloor \leq r(BRC^{m+n}) \leq 2m + 2n$.

4 Simplex DNA code of type α and type β over \mathbb{N}

In ref. [3] has been studied of Quaternary simplex codes of type α and type β . Type α Simplex code S_k^α is a linear dna code over \mathbb{N} with parameters $[4^k, k]$ and an inductive generator matrix given by

$$GM_k^\alpha = \left[\begin{array}{c|c|c|c} A \cdots A & C \cdots C & T \cdots T & G \cdots G \\ \hline GM_{k-1}^\alpha & GM_{k-1}^\alpha & GM_{k-1}^\alpha & GM_{k-1}^\alpha \end{array} \right] \tag{3}$$

with $GM_1^\alpha = [A(0) C(1) T(2) G(3)]$. Type simplex code S_k^β is a punctured version of S_k^α with parameters $[2^{k-1}, (2^k - 1), k]$ and an inductive generator matrix given by

$$GM_2^\beta = \left[\begin{array}{c|c|c} C C C C & A & T \\ \hline A C T G & C & C \end{array} \right] \tag{4}$$

$$GM_k^\beta = \left[\begin{array}{c|c|c} CC \cdots C & AA \cdots A & TT \cdots T \\ \hline GM_{k-1}^\alpha & GM_{k-1}^\beta & GM_{k-1}^\beta \end{array} \right] \tag{5}$$

and for $k > 2$, where GM_{k-1}^α is the generator matrix of S_{k-1}^α . For details the reader is referred to [3]. Type α code with minimum chinese euclidean weight is 8.

Theorem 4.1. $r(S_k^\alpha) \leq 2^{2k+1} - 3$.

Proof. Let $x = CC \cdots C \in \mathbb{N}^n$. By equation(3) the result of Mattson for finite rings and using Theorem 3.2, then

$$\begin{aligned} r(S_k^\alpha) &\leq r(S_{k-1}^\alpha) + r(\langle \overbrace{CC \cdots C}^{2^{2(k-1)}} \overbrace{TT \cdots T}^{2^{2(k-1)}} \overbrace{GG \cdots G}^{2^{2(k-1)}} \rangle) \\ &= r(S_{k-1}^\alpha) + 6 \cdot 2^{2(k-1)} \\ &= 6 \cdot 2^{2(k-1)} + 6 \cdot 2^{2(k-2)} + 6 \cdot 2^{2(k-3)} + \dots + 6 \cdot 2^{2 \cdot 1} + r(S_1^\alpha) \\ r(S_k^\alpha) &\leq 2^{2k+1} - 3 \text{ (since } r(S_1^\alpha) = 5 \text{)}. \end{aligned}$$

□

Theorem 4.2. $r(S_k^\beta) \leq 2^k(2^k - 1) - 7$

Proof. By equation(5), Proposition 3 and Theorem 3.4, thus

$$\begin{aligned} r(S_k^\beta) &\leq r(S_{k-1}^\beta) + r(\langle \overbrace{CC \cdots C}^{4^{(k-1)}} \overbrace{TT \cdots T}^{2^{(2k-3)} - 2^{(k-2)}} \rangle) \\ &= r(S_{k-1}^\beta) + 2^{(2k-2)} + 2^{(2k-3)} - 2^{(k-2)} \\ &\leq 2(2^{(2k-2)} + 2^{(2k-4)} + \dots + 2^4) + 2(2^{(2k-3)} + 2^{(2k-5)} + \dots + 2^3) - \\ &\quad 2(2^{(k-2)} + 2^{(k-3)} + \dots + 2) + r(S_2^\beta) \\ r_{CE}(S_k^\beta) &\leq 2^{k-1}(2^k - 1) - 7 \text{ (since } r(S_2^\beta) = 5 \text{)}. \end{aligned}$$

□

5 MacDonald DNA codes of type α and β over \mathbb{N}

The q -ary MacDonald code $M_{k,t}(q)$ over the finite field F_q is a unique

$\left[\frac{q^k - q^t}{q - 1}, k, q^{k-1} - q^{t-1} \right]$ code in which every non-zero codeword has weight either q^{k-1} or $q^{k-1} - q^{t-1}$ [10]. In [12], he studied the covering radius of MacDonald codes over a finite field. In fact, he has given many exact values for smaller dimension. In [8], authors have defined the MacDonald codes over a ring using the generator matrices of simplex codes. For $2 \leq t \leq k - 1$, let $GM_{k,t}^\alpha$ be the matrix obtained from GM_k^α by deleting columns corresponding to the columns of GM_t^α . That is,

$$GM_{k,t}^\alpha = \left[GM_k^\alpha \setminus \frac{0}{GM_t^\alpha} \right] \tag{6}$$

and let $GM_{k,t}^\beta$ be the matrix obtained from GM_k^β by deleting columns corresponding to the columns of GM_t^β . That is,

$$GM_{k,t}^\beta = \left[GM_k^\beta \setminus \frac{0}{GM_t^\beta} \right] \tag{7}$$

where $[A \setminus B]$ denotes the matrix obtained from the matrix A by deleting the columns of the matrix B and 0 is a $(k - t) \times 2^{2t} ((k - t) \times 2^{t-1} (2^t - 1))$. The code generated by the matrix $GM_{k,t}^\alpha$ is called code of type α and the code generated by the matrix $GM_{k,t}^\beta$ is called Macdonald code of type β . The type α code is denoted by $M_{k,t}^\alpha$ and the type β code is denoted by $M_{k,t}^\beta$. The $M_{k,t}^\alpha$ code is $[4^k - 4^t, k]$ code over \mathbb{N} and $M_{k,t}^\beta$ is a $[(2^{k-1} - 2^{t-1})(2^k + 2^t - 1), k]$ code over \mathbb{N} . In fact, these codes are punctured code of S_k^α and S_k^β respectively.

Next Theorem gives a basic bound on the covering radius of above Macdonald codes.

Theorem 5.1. $r(M_{k,t}^\alpha) \leq 2^{2k+1} - 2^{2r+1} + r(M_{r,t}^\alpha)$ for $t < r \leq k$.

Proof. In equation(6), Proposition 3 and Theorem 3.2, thus

$$\begin{aligned} r(M_{k,t}^\alpha) &\leq r(\langle \overbrace{CC \cdots C}^{2^{2(k-1)}} \overbrace{TT \cdots T}^{2^{2(k-1)}} \overbrace{GG \cdots G}^{2^{2(k-1)}} \rangle) + r(M_{r,t}^\alpha) \\ &= 6.4^{k-1} + r(M_{k-1,t}^\alpha), \text{ for } k \geq r > t. \\ &\leq 6.4^{k-1} + 6.4^{k-2} + \cdots + 6.4^r + r(M_{r,t}^\alpha) \text{ for } k \geq r > t \\ r(M_{k,t}^\alpha) &\leq 2^{2k+1} - 2^{2r+1} + r(M_{r,t}^\alpha), \text{ for } k \geq r > t. \end{aligned}$$

□

Theorem 5.2. $r(M_{k,t}^\beta) \leq 2^k (2^k - 1) + 2^r (1 - 2^r) + r(M_{r,t}^\beta)$, for $t < r \leq k$.

Proof. Using Proposition 3, Theorem 3.4 and in equation(7), obtain


$$\begin{aligned} r(M_{k,t}^\beta) &\leq r(\langle \overbrace{CC \cdots C}^{2^{2(k-1)} \ 2^{2(k-1)-1} \ \cdots \ 2^{(k-1)-1}} \overbrace{TT \cdots T}^{2^{2(k-1)-1} \ \cdots \ 2^{(k-1)-1}} \rangle) + r(M_{k-1,t}^\beta) \\ &\leq 2.2^{2(k-1)+2.2^{2(k-1)-1} - 2^{(k-1)-1} + r(M_{k-1,t}^\beta) \\ &= 2.2^{2(k-1)+2.2^{2(k-1)-1} - 2^{(k-1)-1} + 2 \cdot 2^{2(k-2)} + 2.2^{2(k-2)-1} - 2.2^{2(k-2)-1} + r_{CE}(M_{k-2,t}^\beta) \\ &\leq 2.2^{2(k-1)+2.2^{2(k-1)-1} - 2^{(k-1)-1} + 2.2^{2(k-2)} + 2.2^{2(k-2)-1} - 2.2^{2(k-2)-1} + \cdots + \\ &2.2^{2r} + 2.2^{2r-1} + 2.2^{r-1} + r(M_{r,t}^\beta) \\ &= 2^{2k} - 2^{2r} - 2^k + 2^r + r(M_{r,t}^\beta), t < r \leq k. \\ r(M_{k,t}^\beta) &\leq 2^k (2^k - 1) + 2^r (1 - 2^r) + r(M_{r,t}^\beta), t < r \leq k. \end{aligned}$$

□

References

- [1] M. L. Adleman, Molecular computation of solutions to combinatorial problems, *Science* 266 (1994) 1021-1024.
- [2] T. Aoki, P. Gaborit, M. Harada, M. Ozeki, P. Sol'e, On the covering radius of Z_4 codes and their lattices, *IEEE Trans. Inform. Theory* 45(6) (1999) 2162-2168.
- [3] M. C. Bhandari, M. K. Gupta, A. K. Lal, On Z_4 simplex codes and their gray images, *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes, AAECC- 13, Lecture Notes in Computer Science* 1719(1999) 170-180.
- [4] A. Bonnetcaze, P. Sol'e, C. Bachoc, B. Murrain, Type II codes over Z_4 , *IEEE Trans. Inform. Theory* 43 (1997) 969-976.
- [5] P. Chella Pandian, On the covering radius of codes over Z_4 with chinese euclidean weight, *International Journal on Information Theory (IJIT)* 4(4) (2015) 1-8.
- [6] G. D. Cohen, M. G. Karpovsky, H. F. Mattson, J. R. Schatz, Covering radius- survey and recent results, *IEEE Trans. Inform. Theory* 31(3) (1985) 328-343.
- [7] C. Cohen, A. Lobstein, N. J. A. Sloane, Further results on the covering radius of codes, *IEEE Trans. Inform. Theory* 32(5) (1986) 680-694.
- [8] C. J. Colbourn, M. K. Gupta, On quaternary MacDonal codes, in: *Proc. Information Technology: Coding and Computing (ITCC), 2003*, 212-215.
- [9] J. H. Conway, N. J. A. Sloane, Self-dual codes over the integers modulo 4, *Journal of Combin. Theory Ser. A* 62 (1993) 30-45.
- [10] S. Dodunekov, J. Simonis, Codes and projective multisets. *Electron. J. Combin* 5(1) (1998) 23.
- [11] S. T. Dougherty, M. Harada, P. Sol'e, Shadow codes over Z_4 , *Finite Fields and Their Appl.* (to appear).
- [12] C. Durairajan, On Covering Codes and Covering Radius of Some Optimal Codes, Ph. D. Thesis, Department of Mathematics, IIT Kanpur, 1996.
- [13] M. K. Gupta, C. Durairajan, On the covering radius of some modular codes, *Journal of Advances in Mathematics of Computations* 8(2) (2014) 9.
- [14] M. K. Gupta, G. Glynn David, T. Aaron Gulliver, On Senary Simplex Codes, In: S. Boztaş, I.E. Shparlinski, (eds) *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes. AAECC. Lecture Notes in Computer Science, 2227*. Springer, Berlin, Heidelberg, 2001.
- [15] A. R. Hammons, P. V. Kumar, A. R. Calderbank, N. J. A. Sloane, P., Sol'e, The Z_4 -linearity of kerdock, preparata, goethals, and related codes, *IEEE Trans. Inform. Theory* 40(1994) 301-319.
- [16] M. Harada, New extremal Type II codes over Z_4 , *Des. Codes and Cryptogr.* 13 (1998) 271-284 .
- [17] D. J. Watson, C. H. F. Crick, A structure for deoxyribose nucleic acid, *Nature.* 25 (1953) 737-738.

Citation: P. Chella Pandian, On the Covering Radius of DNA Code over a Finite Ring, *J. Disc. Math. Appl.* 8(2) (2023) 73–80.

 <https://doi.org/10.22061/jdma.2023.1943>



COPYRIGHTS

©2023 The author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.